



## King's Research Portal

DOI:

[10.1016/j.ydbio.2009.09.025](https://doi.org/10.1016/j.ydbio.2009.09.025)

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Christophorou, N. A. D., Bailey, A. P., Hanson, S., & Streit, A. (2009). Activation of Six1 target genes is required for sensory placode formation. *Developmental Biology*, 336(2), 327 - 336.

<https://doi.org/10.1016/j.ydbio.2009.09.025>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.


### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



## Genomes &amp; Developmental Control

## Activation of Six1 target genes is required for sensory placode formation

Nicolas A.D. Christophorou, Andrew P. Bailey<sup>1</sup>, Sven Hanson, Andrea Streit<sup>\*</sup>

Department of Craniofacial Development, King's College London, Guy's Campus, Tower Wing Floor 27, London SE1 8RT, UK

## ARTICLE INFO

## Article history:

Received for publication 12 June 2009

Revised 16 September 2009

Accepted 16 September 2009

Available online 23 September 2009

## Keywords:

Chick

Eyes absent

Ear

Eye

Lens

Olfactory epithelium

Pax2

Pax3

Pax6

Sine oculis

## ABSTRACT

In vertebrates, cranial placodes form crucial parts of the sensory nervous system in the head. All cranial placodes arise from a common territory, the preplacodal region, and are identified by the expression of Six1/4 and Eya1/2 genes, which control different aspects of sensory development in invertebrates as well as vertebrates. While So and Eya can induce ectopic eyes in *Drosophila*, the ability of their vertebrate homologues to induce placodes in non-placodal ectoderm has not been explored. Here we show that Six1 and Eya2 are involved in ectodermal patterning and cooperate to induce preplacodal gene expression, while repressing neural plate and neural crest fates. However, they are not sufficient to induce ectopic sensory placodes in future epidermis. Activation of Six1 target genes is required for expression of preplacodal genes, for normal placode morphology and for placode-specific Pax protein expression. These findings suggest that unlike in the fly where the Pax6 homologue Eyeless acts upstream of Six and Eya, the regulatory relationships between these genes are reversed in early vertebrate placode development.

© 2009 Elsevier Inc. All rights reserved.

## Introduction

In *Drosophila*, *sine oculis* (*so*) and *eyes absent* (*eya*) are nuclear factors that play a key role during compound eye development. Together with seven other transcriptional regulators they form the retinal determination (RD) network, a complex gene regulatory network that controls photoreceptor cell specification in the eye-antennal disc (reviewed in Treisman, 1999; Kumar and Moses, 2001; Donner and Maas, 2004; Pappu and Mardon, 2004). These genes have been placed into a functional network because of their overlapping expression patterns (Bessa et al., 2002) as well as results of loss- and gain-of-function experiments (Bonini et al., 1993; Cheyette et al., 1994; Mardon et al., 1994; Quiring et al., 1994; Serikaku and O'Tousa, 1994; Halder et al., 1995; Bonini et al., 1997; Shen and Mardon, 1997; Czerny et al., 1999; Seimiya and Gehring, 2000; Weasner et al., 2007) and biochemical data (Chen et al., 1997; Pignoni et al., 1997; Czerny et al., 1999; Niimi et al., 1999; Punzo et al., 2002; Ostrin et al., 2006). Like other members of the RD network, So and Eya function is required for normal eye formation and they have the unique ability to induce ectopic eyes when misexpressed in non-retinal tissue and appear to act synergistically (Bonini et al., 1997; Pignoni et al., 1997; Seimiya and Gehring, 2000; Salzer and Kumar, 2009).

In the eye-antennal disc, *so* and *eya* are downstream of Eyeless (Ey) and Twin of Eyeless (Toy) (Halder et al., 1998; Niimi et al., 1999; Bui et al., 2000; Punzo et al., 2002); they act as a complex to activate downstream target genes and positively feed back on *ey* expression itself (Pignoni et al., 1997). As in the fly, vertebrate homologues of the RD network play an important role in eye development but in addition control different aspects of ear, olfactory and sensory ganglia formation including neurogenesis and proliferation (reviewed in Kawakami et al., 2000; Wawersik and Maas, 2000; Hanson, 2001; Donner and Maas, 2004; Silver and Rebay, 2005; Kumar, 2009). They are therefore considered to be key players in controlling cell fate determination in the cranial sensory nervous system, although their regulatory relationship is not always similar to that described in *Drosophila* (for detailed discussion see: Donner and Maas, 2004; Kumar, 2009). In vertebrates, the cranial sensory nervous system largely arises from specialised epithelia, the sensory placodes (reviewed in Baker and Bronner-Fraser, 2001; Streit, 2004; Schlosser, 2006; Streit, 2007). Even before these become morphologically distinct, *so* and *eya* homologues (Six1 and -4; Eya1 and -2) identify most, if not all, placode progenitor cells in what has been called the preplacodal region (PPR) located next to the anterior neural plate (Mishima and Tomarev, 1998; Esteve and Bovolenta, 1999; Sahly et al., 1999; Kobayashi et al., 2000; Pandur and Moody, 2000; Streit, 2002; McLarren et al., 2003; Bessarab et al., 2004; Bhattacharyya et al., 2004; Schlosser and Ahrens, 2004; Litsiou et al., 2005; Xu et al., 2008). Interestingly, their expression begins before that of the *ey* homolog Pax6 or of any other member of the Pax gene family later found in

<sup>\*</sup> Corresponding author.E-mail address: [andrea.streit@kcl.ac.uk](mailto:andrea.streit@kcl.ac.uk) (A. Streit).<sup>1</sup> Present address: NIMR, Developmental Neurobiology, Mill Hill, London NW7 1AA, UK.

different placodes (Pax2: epibranchial and otic; Pax3: trigeminal; Pax6: lens and olfactory; Li et al., 1994; Stark et al., 1997; Groves and Bronner-Fraser, 2000; Bhattacharyya et al., 2004). These observations suggest that Six and Eya proteins may play a crucial role during early sensory progenitor specification and, unlike in *Drosophila*, may act upstream of Pax genes.

Indeed, a recent study in *Xenopus* revealed that Six1 function is required for cells to acquire preplacodal character and that its misexpression leads to upregulation of genes specific for placode precursors at the expense of neural crest and epidermis (Brugmann et al., 2004). Likewise, in Six1 and -4 compound mutant mice the olfactory placode does not form (Chen et al., 2009), consistent with an early, synergistic function of both genes before placode formation. However, it has not yet been tested whether Six and Eya proteins are at the top of the genetic cascade that controls sensory fates by regulating the onset of placode-specific Pax gene expression or whether they can induce ectopic placodes, as may be expected from their eye-inducing ability in the fly. The Six family member Six3 is expressed early in the lens territory, regulates Pax6 (Liu et al., 2006) and has been shown to induce ectopic lens-like structures in fish (Oliver et al., 1996). However, this lens-inducing ability is confined to the PPR suggesting that this territory possesses special properties distinct from the remaining non-neural ectoderm.

In support of this idea, recent evidence shows that the acquisition of preplacodal character is an essential step for placode induction: only preplacodal cells are competent to form placodes in response to the appropriate inducing signals (Martin and Groves, 2006). Furthermore, all cells within the placode territory have a common developmental potential: irrespective of their later fate they are initially specified as lens and lens formation must be suppressed for other neurogenic placodes to develop (Bailey et al., 2006). Together, these observations suggest that the PPR has unique properties and that Six and Eya proteins may play an important role for the acquisition of PPR character.

Here we address the question of whether Six1 and Eya2 are sufficient to confer PPR properties to non-placodal cells, whether they are sufficient to induce ectopic placodes and whether they act upstream of placode-specific Pax gene expression. We show that in chick, as in the fly, Six1 and Eya2 act synergistically: together they promote preplacodal gene expression while suppressing neural and neural crest cell fates. While activation of Six1 target genes is required for the specification of placode progenitors, Six1 and Eya2 are not sufficient to impart preplacodal properties (placode competence or lens specification) to cells that normally do not contribute to the sensory nervous system. Likewise, combined expression of Six1 and Eya2 does not induce ectopic placodes. However, unlike in the fly, Six1 appears to act upstream of placode-specific Pax gene expression: activation of Six1 target genes is required for their expression and for ectodermal cells to acquire placode morphology.

## Materials and methods

### Expression constructs and morpholinos

The coding region of human Six1 (Boucher et al., 1996) and chick Eya2 (Mishima and Tomarev, 1998) were cloned into the pCAB-IRES-GFP (Niwa et al., 1991). Engrailed-Six1HD in pCS2 was a kind gift from Dr Sally Moody (Brugmann et al., 2004). Fluorescein-coupled morpholinos leading to deletion of exon3 or exon6 of chick Eya2 and respective control morpholinos were described previously (Mende et al., 2008).

### Embryo culture, electroporation and explant cultures

Fertile hens' eggs (Henry Stewart) were incubated at 38 °C to reach stage HH3<sup>+</sup>/4 or 5–6 (Hamburger and Hamilton, 1951) and cultured

according to a modified version of New's method (New, 1955; Stern and Ireland, 1981). Electroporation of expression constructs at HH3<sup>+</sup>/4 and morpholinos at HH5–6 was conducted as previously described (McLarren et al., 2003; Mende et al., 2008). After overnight culture, embryos were fixed in 4% paraformaldehyde (PFA) in phosphate buffered saline (PBS) for 15 min for immunohistochemistry or overnight at 4 °C for in situ hybridisation.

To assay the ability of Six1 and Eya2 to confer lens specification and FGF2 responsiveness to non-placodal cells, electroporated area pellucida or area opaca epiblast visualised by GFP fluorescence was dissected after 12 h using tungsten needles. The placode territory without underlying mesoderm and endoderm was dissected from stage HH6 embryos as positive control. Explants were kept in Tyrode's saline on ice, until collagen cultures were set up as previously described (Bailey et al., 2006). Explants were grown in vitro at 37 °C in the presence of FGF2 (250 ng/ml; R&D systems; Martin and Groves, 2006) for 20 h to assay otic marker expression or absence of any growth factors for 60–72 h to assay lens specification. Explants were then fixed for 15 min in 4% PFA in PBS for immunohistochemistry.

### Whole-mount in situ hybridisation and immunocytochemistry

Whole-mount in situ hybridisation was performed using digoxigenin (DIG)-labelled antisense RNA probes as previously described (Streit et al., 1998). The following plasmids were used to generate DIG-labelled antisense riboprobes: Dlx5 (McLarren et al., 2003), Eya2 (Mishima and Tomarev, 1998), Gata3 (Sheng and Stern, 1999), GnRH1 (a gift from Dr Ian Dunn), Pax2 (a gift from Dr Martyn Goulding), Pax6 (Li et al., 1994), Pax7 (Basch et al., 2006), Raldh3 (Blentic et al., 2003), Six1 (a gift from Dr Guillermo Oliver), Six4 (Esteve and Bovolenta, 1999) and Sox2 and Sox3 (Uwanogho et al., 1995).

Immunocytochemistry for GFP was performed using a polyclonal anti-GFP antibody (Molecular Probes; 1:2000 in PBS containing 5% sheep serum, 1% Triton X-100) followed by an HRP-conjugated goat anti-rabbit IgG (Jackson ImmunoResearch; 1:500 in PBS containing 5% sheep serum, 1% Triton X-100). Samples were then processed for cryosectioning.

Immunocytochemistry on cryosections was performed using polyclonal antibodies against chick  $\delta$ -crystallin (a gift from Dr J. Piatigorsky), GFP (Molecular probes) and chick Pax2 (Zymed), and monoclonal antibodies against phospho-Histone H3 (Imgenex), Pax3 and Pax6 (Developmental Hybridoma Bank). The appropriate Alexa-fluor 488- and 594-coupled secondary antibodies were purchased from Molecular Probes; nuclei were stained by DAPI (Molecular Probes).

To quantify the number of morpholino or GFP carrying cells expressing Pax2, Pax6 and Pax3 cells in each embryo, digital images from each section were taken after immunostaining using a Leica TCS SP5 confocal microscope. For each section, the total number of morpholino or GFP carrying cells was determined by counting the green fluorescent cells with visible nuclei (MO or GFP/DAPI<sup>+</sup>). The number of Pax2<sup>+</sup>, Pax3<sup>+</sup> and Pax6<sup>+</sup> cells among the morpholino carrying cells was determined by counting the MO or GFP/Pax2, -3 or -6/DAPI<sup>+</sup> cells. An unpaired *t*-test was performed to determine the statistical significance between splice-blocking morpholinos and control morpholinos and between engrailed-Six1HD and GFP electroporated cells.

## Results

### *Six1 and Eya2 promote preplacodal gene expression but are not sufficient to induce ectopic placodes*

To assess whether Six and Eya proteins alone or in combination are sufficient to induce ectopic sensory placodes, as may be expected from their eye-inducing ability in *Drosophila*, or to impart preplacodal




character we used gain-of-function experiments in the chick embryo. After electroporation of primitive streak stage ( $\text{HH3}^+/4$ ) embryos with full length Six1 or Eya2, no changes were observed in markers specific for the neural plate (*Sox2*: Six1  $n=7$ , Eya2  $n=6$ ), neural crest cells (*Pax7*: Six1  $n=7$ , Eya2:  $n=5$ ), non-neural ectoderm (*Dlx5*: Six1  $n=11$ , Eya2:  $n=7$ ; *Gata3*: Six1  $n=7$ ) or the placode territory itself (*Six4*: Six1  $n=11$ , Eya2:  $n=8$ ; *Eya2*: Six1  $n=10$ ) (data not shown). However, when misexpressed together, Six1 and Eya2, but not GFP (controls *Sox2*  $n=8$ ; *Sox3*  $n=7$ ; *Pax7*  $n=7$ ; *Six4*  $n=11$ ; *Dlx5*  $n=11$ ; *Gata3*  $n=5$ ; Figs. 1A–C, A'–C', a–c, G–I, G'–I', g–i), repress *Sox2* and *Sox3* (13/16, 8/11; Figs. 1D, D', d, E, E', e) and *Pax7* (16/27; Figs. 1F, F', f), while inducing ectopic expression of the PPR marker *Six4* in the future epidermis and in the extraembryonic region (13/20; Figs. 1J,


J', j). In addition, misexpression of Six1 and Eya2 leads to non-cell autonomous upregulation of *Dlx5* (7/9; Figs. 1K, K', k, arrow heads) and *Gata3* (6/10; Figs. 1L, L', l, arrow head).

To determine whether Six1 and Eya2 are sufficient to induce ectopic placodes, as might be expected from their eye-inducing ability in the fly, we tested for ectopic expression of placode-specific markers after Six1 and Eya2 misexpression. We never observed ectopic expression of *Pax2* (otic and epibranchial;  $n=9$ ; Figs. 2A, A'), *Pax6* (lens;  $n=14$ ; Figs. 2B, B') or *GnRH1* and *Raldh3* (olfactory;  $n=5$  and  $n=6$ ; Figs. 2C, C', D, D').

It has previously been suggested that high levels of Six1 and Eya1 expression promote proliferation, while low levels promote differentiation into otic neurons (Schlosser et al., 2008). To test whether



**Fig. 1.** Six1 and Eya2 promote preplacodal properties. GFP (A–C, A'–C', G–I, G'–I', a–c, g–i) or Six1 and Eya2 (D–F, D'–F', d–f, J–L, J'–L', j–l) were electroporated into primitive streak stage embryos. After overnight culture, the expression of neural, neural crest, preplacodal and non-neural ectoderm markers was assessed by in situ hybridisation (blue) followed by GFP immunohistochemistry (brown) to visualise transgene carrying cells. Panels A–L show embryos before GFP staining and panels A'–L' the same embryos thereafter. Six1/Eya2 misexpression suppresses *Sox2* (D, D', d), *Sox3* (E, E', e) and *Pax7* (F, F', f) while upregulating *Six4* (J, J', j), *Dlx5* (K, K', k) and *Gata3* (L, L', l). GFP misexpression does not affect the expression of any marker (A–C, A'–C', G–I, G'–I', a–c, g–i). Arrow heads in a–f and j point to electroporated cells (brown), arrow heads in k and l to cells that non-autonomously upregulate *Dlx5* and *Gata3* expression (blue), respectively. Lines in A'–L' correspond to the level of the sections in a'–l'.



**Fig. 2.** Six1 and Eya2 do not promote ectopic placode formation. Six1 and Eya2 were electroporated into primitive streak stage embryos. After 20–24 h, the expression of otic (A, A'), lens (B, B') and olfactory markers (C, C', D, D') was assessed by in situ hybridisation (blue; A–D). Electroporated cells were visualised by GFP immunohistochemistry (brown; A'–D'). Upregulation of Pax2 (A, A'), Pax6 (B, B'), Gnrh (C, C') and Raldh3 (D, D') was never observed.

Six1 and Eya2 increase proliferation in epiblast cells we compared phospho-Histone H3 staining in GFP (control;  $n = 7$ ) and Six1/Eya2 ( $n = 7$ ) electroporated embryos. No difference was observed (data not shown) suggesting that the failure to differentiate into placodes is not due to increased proliferation of cells carrying exogenous Six1 and Eya2.

These results suggest that while Six1 and Eya2 act together to promote preplacodal character, they cannot induce placode-specific gene expression in non-placodal ectoderm.

#### *Six1 and Eya2 do not account for all PPR properties*

The PPR is a unique part of the neurula stage ectoderm: it is the only region responsive to otic inducing signals and it is initially specified as lens (Bailey et al., 2006; Martin and Groves, 2006). We therefore assessed whether Six1 and Eya2 are sufficient to confer these properties to cells not fated to become placodes. Six1 and Eya2 were misexpressed at the border of the embryonic and extraembryonic region at gastrula stages (HH3<sup>+</sup>/4). After overnight culture, the electroporated tissue was excised and cultured in the presence or absence of the otic inducing signal FGF2 for 20–24 h (Fig. 3A). PPR from HH6 stage embryos was used as a positive control. While 87.5% of PPR explants expressed the otic marker Pax2 (7/8; Figs. 3B–D), neither FGF2 treated ( $n = 14$ ; Figs. 3E–G) nor untreated ( $n = 11$ ; data not shown) tissue carrying exogenous Six1 and Eya2 did. Likewise, unlike endogenous PPR from HH6 (8/9; Figs. 3H–J), Six1 and Eya2 electroporated epiblast cultured for 60–72 h in the absence of growth factors does not acquire lens character as assessed by  $\delta$ -crystallin expression ( $n = 15$ ; Figs. 3K–M). These results show that although Six1 and Eya2 can induce PPR markers in non-placodal ectoderm (see above), together they are not sufficient to impart PPR properties to cells not normally fated to become placodes.

#### *Loss of Eya2 causes a moderate decrease of Pax2 expression*

While Six1 and Six4 are initially weakly expressed in the neural plate before becoming confined to the PPR, Eya2 is confined to the PPR from the onset of its expression. Although weak expression of Eya1 has been reported in the anterior PPR (Ishihara et al., 2008), in our hands this transcript cannot be detected. We therefore asked whether Eya2 is required for PPR specification and placode formation. Two different morpholinos were electroporated into placode precursors at

primitive streak to head process stages (HH3<sup>+</sup>–5) and the expression of PPR and placode markers was assessed after 15–24 h culture. No change in Six1 or Six4 expression was observed (Six1  $n = 10$ ; Six4  $n = 8$ ; data not shown), and Eya2 MO ( $n = 13$ ) carrying cells appear to integrate into the otic placode to the same extent as control MO electroporated cells ( $n = 9$  for each). However, a very moderate reduction of Pax2 protein in the otic placode is observed in Eya2 MO carrying cells when compared to controls (control:  $99.66\% \pm 0.34$ , Eya2 MO  $89.33\% \pm 1.19$  of all electroporated cells are Pax2<sup>+</sup>,  $p = 0.021$ ; Supplementary Fig. 1). When Eya2 MOs are targeted to the lens territory (Eya2 MO  $n = 9$ ; control MO  $n = 10$ ) no changes in the early lens marker Pax6 are observed.


We next investigated whether Eya2 is required for PPR properties, such as competence to respond to otic inducing signals or lens specification. Eya2 or control MOs were electroporated into placode precursors, the targeted tissue was excised and cultured overnight in the presence or absence of FGF2 and assayed for the expression of the otic marker Pax2 after FGF2 treatment and for the early lens marker Pax6. There is no difference in Pax2 induction by FGF2 or in the number of Pax6<sup>+</sup> (i.e., lens specified) cells between Eya2 and control MO electroporated explants.

In summary, while Eya2 knock-down leads to a moderate reduction in expression of the otic marker Pax2, PPR character remains unchanged when Eya2 is reduced.

#### *Six1 regulates gene expression at the neural plate border*

Unlike Eya proteins, Six1 directly binds to specific DNA target sequences via its homeodomain and acts as an activator in association with Eya proteins (Li et al., 2003; Rayapureddi et al., 2003; Tootle et al., 2003; Hu et al., 2008). Our results above show that misexpression of Six1 and Eya2 leads to upregulation of PPR specific gene expression (Fig. 1). To investigate whether activation of Six1 target genes is required, we used an expression construct where the homeodomain of Six1 is fused to a constitutive repressor (EnR). Electroporation of EnR-Six1, but not of GFP (Figs. 4E–E', F–F') at gastrula stages (HH3<sup>+</sup>/4) leads to loss of the preplacodal markers Six4 (8/10; Figs. 4A–A', a) and Eya2 (4/8; Figs. 4B–B', b), while Dlx5 expression in the placode territory remains unchanged ( $n = 6$ ; data not shown). Occasionally, we observe an expansion of the neural marker Sox2 into the non-neural ectoderm (6/11; Figs. 4C–C', c) suggesting that when Six1 target genes are repressed, cells at the neural plate border can acquire





**Fig. 3.** Six1 and Eya are not sufficient to confer preplacodal properties to non-placodal cells. (A) To assess whether Six1 and Eya2 impart responsiveness to otic inducing signals or lens specification to cells that normally do not form sensory placodes, Six1 and Eya2 were electroporated into the ectoderm at HH3<sup>+</sup>/4; embryos were grown until they had reached HH5/6 and the electroporated cells were isolated and cultured in the presence of FGF2 for 24 h to assess otic markers or in the absence of growth factors for 2.5 days to assess lens markers. PPR from HH5/6 embryos was used as positive control. In the presence of FGF2, PPR explants (B–D) express the otic marker Pax2 (red in C, magenta in D), while Six1/Eya2 electroporated ectoderm does not (E–G; green). After 2.5 days in culture (H–J), PPR explants express lens specific  $\delta$ -crystallin (red in I, J). However, Six1/Eya2 electroporated ectoderm (K–M, green) is negative for  $\delta$ -crystallin. D, G, J and M show nuclei stained with DAPI.

neural character. In contrast, repression of Six1 target genes in the anterior neural folds, which normally do not generate neural crest cells, leads to expansion of the neural crest marker *Pax7* (7/11; Figs. 4D–D', d) and we occasionally observe lateral expansion of *Pax7* into the placode territory.

Thus, repression of Six1 target genes and misexpression of Six1/Eya2 show opposite phenotypes (compare Figs. 1 and 4), suggesting that in the context of sensory precursor specification Six1 acts as an activator. Preplacodal expression of *Six4* and *Eya2* is dependent on Six1 or its targets, while preventing the formation of neural and neural crest cells in tissue that normally forms placodes may depend on activation of a repressor.


#### *Six1 regulates Pax gene expression in placode progenitors and placodal morphology*

In placode precursors, Six and Eya genes are expressed prior to the onset of placode-specific Pax gene expression and prior to the formation of morphological placodes. Our results above show that misexpression of Six1/Eya2 is not sufficient to induce placodes in

non-placodal ectoderm; however, activation of Six1 targets may be required. Indeed, misexpression of EnR-Six1 leads to severe malformation of the otic placode ( $n=13$ ; Figs. 5A b, b', b'', c c'), which becomes split into multiple 'cuplets'. Cells carrying the transgene fail to acquire columnar morphology typical of the placode and do not express Pax2 (70.2%;  $n=7$ ; Figs. 5A b, b', b'', d). EnR-Six1 expressing cells appear to cluster and disrupt the organisation of the otic placode. Likewise, when misexpressed in the trigeminal and lens territory, EnR-Six1 cells fail to express Pax3 (44.15%;  $n=7$ ; Figs. 5B, b, b', b'', c) and Pax6 (49.55%;  $n=7$ ; Figs. 5C, b, b', b'', c), respectively. In contrast, control electroporated embryos never show any placode defects or significant loss of Pax protein expression (Pax2: 9.96%, Figs. 5A a, a', a'', d  $n=6$ ; Pax3: 17.4%, Figs. 5B a, a', a'', c,  $n=4$ ; Pax6: 5.4%,  $n=7$ , Figs. 5C a, a', a'', c). Statistical analysis shows that the differences observed between EnR-Six1 and GFP misexpression are significant (Pax2:  $p=0.0000016$ , Fig. 5A d; Pax3:  $p=0.0003$ , Fig. 5B c; Pax6:  $p=0.0000027$ , Fig. 5C c). Thus, activation of Six1 target genes is required for the expression of placode-specific Pax proteins. However, whether Six proteins directly activate Pax gene transcription or whether intermediate factors are required remains to be







**Fig. 5.** Activation of Six1 target genes is required for placode-specific Pax protein expression and placode formation. EnR-Six1 or GFP was electroporated into the future otic (A), trigeminal (B) or lens (C) region; at HH9<sup>+</sup>–11 the expression of Pax2 (A), Pax3 (B) and Pax6 (C) was assessed by immunocytochemistry in transverse sections. (A) In the otic placode, cells misexpressing GFP (white arrow heads in a–a'', green) continue to be Pax2<sup>+</sup> (red in a', magenta in a''). However, in the presence of EnR-Six1 (white arrow heads in b–b'', green), Pax2 expression is lost (red in b', magenta in b'') and otic placode morphology is disrupted. White brackets indicate the formation of multiple small otic cups. (C, c') Dorsal view (anterior towards the left) of an embryo electroporated with EnR-Six1 into the left otic region (green) and stained for Pax2 (magenta). C' shows a higher magnification of the area indicated by the white square in c. Note the disruption of otic morphology and formation of multiple smaller cups. To quantify the effect, the percentage of Pax2<sup>+</sup> cells among electroporated cells was counted in 7 EnR-Six1 and 6 GFP expressing embryos (d). Pax2 expression is reduced to approximately 30% when Six1 target gene activation is inhibited ( $p = 1.6 \times 10^{-6}$ ). (B) In the trigeminal placode, cells misexpressing GFP (white arrow heads in a–a'', green) continue to be Pax3<sup>+</sup> (red in a', magenta in a''). However, in the presence of EnR-Six1 (white arrow heads in b–b'', green), Pax3 expression is lost (red in b', magenta in b''). The effect was quantified by counting Pax3<sup>+</sup> cells among electroporated cells in the trigeminal region in 7 EnR-Six1 and 4 GFP expressing embryos (c). Pax3 expression is reduced to approximately 50% when Six1 target gene activation is inhibited ( $p = 3.0 \times 10^{-4}$ ). (C) In the lens placode, cells misexpressing GFP (white arrow heads in a–a'', green) continue to be Pax6<sup>+</sup> (red in a', magenta in a''). However, in the presence of EnR-Six1 (white arrow heads in b–b'', green), Pax6 expression is lost (red in b', magenta in b''). The effect was quantified by counting Pax6<sup>+</sup> cells in the lens placode in 7 EnR-Six1 and 7 GFP expressing embryos (c). Pax6 expression is reduced to approximately 50% when Six1 target gene activation is inhibited ( $p = 2.7 \times 10^{-6}$ ).



One surprising difference between previous *Xenopus* and chick results is that *Six1* can act alone to pattern the ectoderm in *Xenopus*, while in chick misexpression of *Six1* together with the co-activator *Eya2* is required. The latter is reminiscent to the situation in *Drosophila*, where the eye-inducing activity of *So* and *Eya* alone is relatively low, but increases considerably when both are co-expressed in non-retinal tissue (Bonini et al., 1997; Pignoni et al., 1997; Seimiya and Gehring, 2000; Salzer and Kumar, 2009). The main differences between the *Xenopus* and chick experiments are timing and the tissue expressing exogenous constructs: while in *Xenopus* *Six1* was injected into ventro-lateral blastomeres at the 32-cell stage, misexpression experiments in chick were performed at late gastrula stages, shortly before *Six1* and *Eya2* become normally expressed. Frog ventro-lateral blastomeres not only contribute to neural crest and placode cells, but also to mesodermal and endodermal derivatives, which come to underlie the placode territory (Moody, 1987). It has previously been shown that mesodermal signals are required for the induction of the placode territory (Litsiou et al., 2005). It is therefore possible that changes in the mesoderm, where the *Six*/*Eya* network participates in somite formation (Heanue et al., 1999), indirectly influence the overlying ectoderm. Another possibility is that *Eya1*, which is already expressed at early gastrula stages in *Xenopus* (David et al., 2001), may interact with exogenous *Six1* to affect patterning of the ectoderm, whereas *Eya1* and *-2* are absent at this stage in the chick (McLarren et al., 2003; Litsiou et al., 2005; Ishihara et al., 2008).

#### Are *Six1* and *Eya2* specifiers for placodal progenitors?

Thus, *Six* and *Eya* proteins appear to be part of a regulatory network of transcription factors that controls cell fate specification at the border of the neural plate and ensures the segregation of neural, neural crest, placodal and epidermal fates. *Six1* and *Eya2* promote preplacodal gene expression, but are they sufficient to endow placode progenitor properties onto cells that normally never contribute to placodes? Recent evidence suggests that to generate sensory placodes, ectodermal cells first have to acquire a 'preplacodal state', which allows them to respond to placode-specific inducing signals (Martin and Groves, 2006). In addition, preplacodal cells initially share a common developmental potential and are initially specified as lens, before lens suppressing and neurogenic placode inducing signals generate placode diversity (Bailey et al., 2006). Here we show that while misexpression of *Six1* and its cofactor *Eya2* can induce preplacodal gene expression, their expression alone does not account for all properties of placode progenitors. Cells expressing exogenous *Six1*/*Eya2* cannot respond to FGF2 to activate otic markers, nor are they lens specified.

Based on the eye-inducing activity of *So* and *Eya* in *Drosophila*, we predicted that *Six1* and *Eya2* may induce ectopic placodes in tissue not normally fated to contribute to the cranial sensory nervous system. However, we find that misexpression of both genes does not lead to ectopic morphological placodes and does not initiate the expression of placode-specific gene expression.

Together, these results suggest that additional factors are required for cells to acquire preplacodal character and for the induction of ectopic placodes in future epidermis. Indeed, in *Drosophila* *Six1* binding partners in addition to *Eya* and Groucho proteins have been identified (Giot et al., 2003); however, their role in vertebrate placode formation has not been investigated. Finally, we cannot exclude the possibility that more subtle regulation of *Six1* and *Eya2* expression levels is required for ectodermal cells to undergo placode differentiation.

#### *Six1* acts upstream of placode-specific *Pax* genes

In *Drosophila* the expression and function of *so* and *eya* in the eye are dependent on the function of the *Pax6* homologues *Ey* and *Toy*.

Surprisingly however, a similar regulatory relationship is not conserved in the vertebrate sensory system. While *Pax6* has retained its activity as a key regulator of vertebrate eye development, its early expression in the future lens ectoderm depends on direct activation by *Six3* via the ectodermal enhancer (Liu et al., 2006). Likewise, in the murine olfactory placode, *Pax6* expression requires *Six1* and *Six4* function (Chen et al., 2009) and the onset of *Six3* and *Eya1* expression is independent of *Pax6* (Purcell et al., 2005). In the otic vesicle, *Pax2*, *Eya1* and *Six1* are expressed in partially overlapping domains, but *Six1* and *Eya1* expression is independent of *Pax2*, while *Eya1* function is required for *Six1* expression (Zheng et al., 2003; Burton et al., 2004). In the preplacodal region, *Six1*, *Six4* and *Eya2* are clearly present prior to the onset of placode-specific expression of *Pax6*, *Pax3* and *Pax2*. Here we show that although ectopic expression of *Six1* and *Eya2* does not result in the induction of *Pax* genes, the activation of *Six1* downstream targets is required for their expression in the lens, trigeminal and otic placode. Likewise, in zebrafish *Six1* morphants lose otic specific expression of *Pax2a* and *Pax2b* (Bricaud and Collazo, 2006). The regulatory elements that control the onset *Pax* gene expression in sensory placodes have not been identified and therefore it remains to be determined whether their activation by *Six* genes is direct or intermediate factors are required. While *Pax* genes are clearly required for sensory organ development in vertebrates, with *Pax6* being essential for lens and olfactory placode formation (Fujiwara et al., 1994; Collinson et al., 2001; Collinson et al., 2003), *Pax3* for trigeminal (Dude et al., 2009) and *Pax2* for otic development (Torres et al., 1996; Burton et al., 2004), these findings suggest that unlike in the fly eye, *Six* and *Eya*, but not *Pax* proteins, are at the top of the genetic hierarchy that specifies sensory precursor cells in vertebrates.

Together our findings propose that while some of the molecular interactions of the RD network are conserved between flies and vertebrates, some of the regulatory relationships have diverged during evolution of the sensory nervous system (for further discussion, see Donner and Maas, 2004; Kumar, 2009). Future studies of sensory placode-specific enhancers will need to clarify the molecular interactions of RD network factors.

#### Acknowledgments

We thank Ewa Kolano for excellent technical assistance and Drs. Claudio Stern and Roberto Mayor for discussions and critical reading of the manuscript. Thanks to Sally Moody for providing the EnR-*Six1* construct and all members of the lab for helpful discussions. This work was supported by BBSRC project grants G20323 and D010659 and the Marie Curie EST project number 504025.

#### Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.ydbio.2009.09.025.

#### References

- Akimenko, M.A., Ekker, M., Wegner, J., Lin, W., Westerfield, M., 1994. Combinatorial expression of three zebrafish genes related to distal-less: part of a homeobox gene code for the head. *J. Neurosci.* 14, 3475–3486.
- Bailey, A.P., Bhattacharyya, S., Bronner-Fraser, M., Streit, A., 2006. Lens specification is the ground state of all sensory placodes, from which FGF promotes olfactory identity. *Dev. Cell* 11, 505–517.
- Baker, C.V., Bronner-Fraser, M., 2001. Vertebrate cranial placodes. I. Embryonic induction. *Dev. Biol.* 232, 1–61.
- Basch, M.L., Bronner-Fraser, M., Garcia-Castro, M.I., 2006. Specification of the neural crest occurs during gastrulation and requires *Pax7*. *Nature* 441, 218–222.
- Beanan, M.J., Sargent, T.D., 2000. Regulation and function of *Dlx3* in vertebrate development. *Dev. Dyn.* 218, 545–553.
- Bessa, J., Gebelein, B., Pichaud, F., Casares, F., Mann, R.S., 2002. Combinatorial control of *Drosophila* eye development by *eyeless*, *homothorax*, and *teashirt*. *Genes Dev.* 16, 2415–2427.

- Bessarab, D.A., Chong, S.W., Korzh, V., 2004. Expression of zebrafish *six1* during sensory organ development and myogenesis. *Dev. Dyn.* 230, 781–786.
- Bhattacharyya, S., Bailey, A.P., Bronner-Fraser, M., Streit, A., 2004. Segregation of lens and olfactory precursors from a common territory: cell sorting and reciprocity of *Dlx5* and *Pax6* expression. *Dev. Biol.* 271, 403–414.
- Blentic, A., Gale, E., Maden, M., 2003. Retinoic acid signalling centres in the avian embryo identified by sites of expression of synthesising and catabolising enzymes. *Dev. Dyn.* 227, 114–127.
- Bonini, N.M., Leiserson, W.M., Benzer, S., 1993. The eyes absent gene: genetic control of cell survival and differentiation in the developing *Drosophila* eye. *Cell* 72, 379–395.
- Bonini, N.M., Bui, Q.T., Gray-Board, G.L., Warrick, J.M., 1997. The *Drosophila* eyes absent gene directs ectopic eye formation in a pathway conserved between flies and vertebrates. *Development* 124, 4819–4826.
- Boucher, C.A., Carey, N., Edwards, Y.H., Siciliano, M.J., Johnson, K.J., 1996. Cloning of the human *SIX1* gene and its assignment to chromosome 14. *Genomics* 33, 140–142.
- Bricaud, O., Collazo, A., 2006. The transcription factor *six1* inhibits neuronal and promotes hair cell fate in the developing zebrafish (*Danio rerio*) inner ear. *J. Neurosci.* 26, 10438–10451.
- Brugmann, S.A., Pandur, P.D., Kenyon, K.L., Pignoni, F., Moody, S.A., 2004. *Six1* promotes a placodal fate within the lateral neurogenic ectoderm by functioning as both a transcriptional activator and repressor. *Development* 131, 5871–5881.
- Bui, Q.T., Zimmerman, J.E., Liu, H., Gray-Board, G.L., Bonini, N.M., 2000. Functional analysis of an eye enhancer of the *Drosophila* eyes absent gene: differential regulation by eye specification genes. *Dev. Biol.* 221, 355–364.
- Burton, C., Cole, L.K., Mulheisen, M., Chang, W., Wu, D.K., 2004. The role of *Pax2* in mouse inner ear development. *Dev. Biol.* 272, 161–175.
- Chen, B., Kim, E.H., Xu, P.X., 2009. Initiation of olfactory placode development and neurogenesis is blocked in mice lacking both *Six1* and *Six4*. *Dev. Biol.* 326, 75–85.
- Chen, R., Amoui, M., Zhang, Z., Mardon, G., 1997. Dachshund and eyes absent proteins form a complex and function synergistically to induce ectopic eye development in *Drosophila* [see comments]. *Cell* 91, 893–903.
- Cheyette, B.N., Green, P.J., Martin, K., Garren, H., Hartenstein, V., Zipursky, S.L., 1994. The *Drosophila* sine oculis locus encodes a homeodomain-containing protein required for the development of the entire visual system. *Neuron* 12, 977–996.
- Collinson, J.M., Quinn, J.C., Buchanan, M.A., Kaufman, M.H., Wedden, S.E., West, J.D., Hill, R.E., 2001. Primary defects in the lens underlie complex anterior segment abnormalities of the *Pax6* heterozygous eye. *Proc. Natl. Acad. Sci. U. S. A.* 98, 9688–9693.
- Collinson, J.M., Quinn, J.C., Hill, R.E., West, J.D., 2003. The roles of *Pax6* in the cornea, retina, and olfactory epithelium of the developing mouse embryo. *Dev. Biol.* 255, 303–312.
- Czerny, T., Halder, G., Kloter, U., Souabni, A., Gehring, W.J., Busslinger, M., 1999. Twin of eyeless, a second *Pax-6* gene of *Drosophila*, acts upstream of eyeless in the control of eye development. *Mol. Cell* 3, 297–307.
- David, R., Ahrens, K., Wedlich, D., Schlosser, G., 2001. *Xenopus* *Eya1* demarcates all neurogenic placodes as well as migrating hypaxial muscle precursors. *Mech. Dev.* 103, 189–192.
- Donner, A.L., Maas, R.L., 2004. Conservation and non-conservation of genetic pathways in eye specification. *Int. J. Dev. Biol.* 48, 743–753.
- Dude, C.M., Kuan, C.Y., Bradshaw, J.R., Greene, N.D., Relaix, F., Stark, M.R., Baker, C.V., 2009. Activation of *Pax3* target genes is necessary but not sufficient for neurogenesis in the ophthalmic trigeminal placode. *Dev. Biol.* 326, 314–326.
- Esteve, P., Bovolenta, P., 1999. *cSix4*, a member of the six gene family of transcription factors, is expressed during placode and somite development. *Mech. Dev.* 85, 161–165.
- Feledy, J.A., Beanan, M.J., Sandoval, J.J., Goodrich, J.S., Lim, J.H., Matsuo-Takasaki, M., Sato, S.M., Sargent, T.D., 1999. Inhibitory patterning of the anterior neural plate in *Xenopus* by homeodomain factors *Dlx3* and *Msx1*. *Dev. Biol.* 212, 455–464.
- Fujiwara, M., Uchida, T., Osumi-Yamashita, N., Eto, K., 1994. Uchida rat (rSey): a new mutant rat with craniofacial abnormalities resembling those of the mouse Sey mutant. *Differentiation* 57, 31–38.
- Groves, A.K., Bronner-Fraser, M., 2000. Competence, specification and commitment in otic placode induction. *Development* 127, 3489–3499.
- Halder, G., Callaerts, P., Gehring, W.J., 1995. Induction of ectopic eyes by targeted expression of the eyeless gene in *Drosophila*. *Science* 267, 1788–1792.
- Halder, G., Callaerts, P., Flister, S., Walldorf, U., Kloter, U., Gehring, W.J., 1998. Eyeless initiates the expression of both sine oculis and eyes absent during *Drosophila* compound eye development. *Development* 125, 2181–2191.
- Hamburger, V., Hamilton, H.L., 1951. A series of normal stages in the development of the chick embryo. *J. Morph.* 88, 49–92.
- Hanson, I.M., 2001. Mammalian homologues of the *Drosophila* eye specification genes. *Semin. Cell Dev. Biol.* 12, 475–484.
- Heanue, T.A., Reshef, R., Davis, R.J., Mardon, G., Oliver, G., Tomarev, S., Lassar, A.B., Tabin, C.J., 1999. Synergistic regulation of vertebrate muscle development by *Dach2*, *Eya2*, and *Six1*, homologs of genes required for *Drosophila* eye formation. *Genes Dev.* 13, 3231–3243.
- Hu, S., Mamedova, A., Hegde, R.S., 2008. DNA-binding and regulation mechanisms of the *SIX* family of retinal determination proteins. *Biochemistry* 47, 3586–3594.
- Ishihara, T., Ikeda, K., Sato, S., Yajima, H., Kawakami, K., 2008. Differential expression of *Eya1* and *Eya2* during chick early embryonic development. *Gene Expr. Patterns* 8, 357–367.
- Kaji, T., Artinger, K.B., 2004. *dlx3b* and *dlx4b* function in the development of Rohon-Beard sensory neurons and trigeminal placode in the zebrafish neurula. *Dev. Biol.* 276, 523–540.
- Kawakami, K., Ohto, H., Ikeda, K., Roeder, R.G., 1996. Structure, function and expression of a murine homeobox protein *arec3*, a homologue of *Drosophila* sine oculis gene product, and implication in development. *Nucleic Acids Res.* 24, 303–310.
- Kawakami, K., Sato, S., Ozaki, H., Ikeda, K., 2000. Six family genes—structure and function as transcription factors and their roles in development. *Bioessays* 22, 616–626.
- Kobayashi, M., Osanai, H., Kawakami, K., Yamamoto, M., 2000. Expression of three zebrafish *Six4* genes in the cranial sensory placodes and the developing somites. *Mech. Dev.* 98, 151–155.
- Kumar, J.P., 2009. The sine oculis homeobox (*SIX*) family of transcription factors as regulators of development and disease. *Cell. Mol. Life Sci.* 66, 565–583.
- Kumar, J.P., Moses, K., 2001. Eye specification in *Drosophila*: perspectives and implications. *Semin. Cell Dev. Biol.* 12, 469–474.
- Li, H.S., Yang, J.M., Jacobson, R.D., Pasko, D., Sundin, O., 1994. *Pax-6* is first expressed in a region of ectoderm anterior to the early neural plate: implications for stepwise determination of the lens. *Dev. Biol.* 162, 181–194.
- Litsiou, A., Hanson, S., Streit, A., 2005. A balance of FGF, Wnt and BMP signalling positions the future placode territory in the head. *Development* 132, 4051–4062.
- Liu, W., Lagutin, O.V., Mende, M., Streit, A., Oliver, G., 2006. *Six3* activation of *Pax6* expression is essential for mammalian lens induction and specification. *Embo J.* 25, 5383–5395.
- Li, X., Oghii, K.A., Zhang, J., Krones, A., Bush, K.T., Glass, C.K., Nigam, S.K., Aggarwal, A.K., Maas, R., Rose, D.W., Rosenfeld, M.G., 2003. *Eya* protein phosphatase activity regulates *Six1*-*Dach*-*Eya* transcriptional effects in mammalian organogenesis. *Nature* 426, 247–254.
- Luo, T., Matsuo-Takasaki, M., Sargent, T.D., 2001. Distinct roles for *Distal-less* genes *Dlx3* and *Dlx5* in regulating ectodermal development in *Xenopus*. *Mol. Reprod. Dev.* 60, 331–337.
- Mardon, G., Solomon, N.M., Rubin, G.M., 1994. dachshund encodes a nuclear protein required for normal eye and leg development in *Drosophila*. *Development* 120, 3473–3486.
- Martin, K., Groves, A.K., 2006. Competence of cranial ectoderm to respond to Fgf signaling suggests a two-step model of otic placode induction. *Development* 133, 877–887.
- McLarren, K.W., Litsiou, A., Streit, A., 2003. *DLX5* positions the neural crest and preplacode region at the border of the neural plate. *Dev. Biol.* 259, 34–47.
- Mende, M., Christophorou, N.A., Streit, A., 2008. Specific and effective gene knock-down in early chick embryos using morpholinos but not *prFRPNAi* vectors. *Mech. Dev.* 125, 947–962.
- Mishima, N., Tomarev, S., 1998. Chicken eyes absent 2 gene: isolation and expression pattern during development. *Int. J. Dev. Biol.* 42, 1109–1115.
- Moody, S.A., 1987. Fates of the blastomeres of the 32-cell-stage *Xenopus* embryo. *Dev. Biol.* 122, 300–319.
- New, D.A.T., 1955. A new technique for the cultivation of the chick embryo in vitro. *J. Embryol. Exp. Morph.* 3, 326–331.
- Niimi, T., Seimiya, M., Kloter, U., Flister, S., Gehring, W.J., 1999. Direct regulatory interaction of the eyeless protein with an eye-specific enhancer in the sine oculis gene during eye induction in *Drosophila*. *Development* 126, 2253–2260.
- Niwa, H., Yamamura, K., Miyazaki, J., 1991. Efficient selection for high-expression transfectants with a novel eukaryotic vector. *Gene* 108, 193–199.
- Oliver, G., Loosli, F., Koster, R., Wittbrodt, J., Gruss, P., 1996. Ectopic lens induction in fish in response to the murine homeobox gene *Six3*. *Mech. Dev.* 60, 233–239.
- Ostrin, E.J., Li, Y., Hoffman, K., Liu, J., Wang, K., Zhang, L., Mardon, G., Chen, R., 2006. Genome-wide identification of direct targets of the *Drosophila* retinal determination protein Eyeless. *Genome Res.* 16, 466–476.
- Ozaki, H., Nakamura, K., Funahashi, J., Ikeda, K., Yamada, G., Tokano, H., Okamura, H.O., Kitamura, K., Muto, S., Kotaki, H., Sudo, K., Horai, R., Iwakura, Y., Kawakami, K., 2004. *Six1* controls patterning of the mouse otic vesicle. *Development* 131, 551–562.
- Pandur, P.D., Moody, S.A., 2000. *Xenopus* *Six1* gene is expressed in neurogenic cranial placodes and maintained in the differentiating lateral lines. *Mech. Dev.* 96, 253–257.
- Pappu, K.S., Mardon, G., 2004. Genetic control of retinal specification and determination in *Drosophila*. *Int. J. Dev. Biol.* 48, 913–924.
- Pera, E., Stein, S., Kessel, M., 1999. Ectodermal patterning in the avian embryo: epidermis versus neural plate. *Development* 126, 63–73.
- Pignoni, F., Hu, B., Zavitz, K.H., Xiao, J., Garrity, P.A., Zipursky, S.L., 1997. The eye-specification proteins *So* and *Eya* form a complex and regulate multiple steps in *Drosophila* eye development [published erratum appears in *Cell* 1998 Feb 20;92 (4):following 585]. *Cell* 91, 881–891.
- Punzo, C., Seimiya, M., Flister, S., Gehring, W.J., Plaza, S., 2002. Differential interactions of eyeless and twin of eyeless with the sine oculis enhancer. *Development* 129, 625–634.
- Purcell, P., Oliver, G., Mardon, G., Donner, A.L., Maas, R.L., 2005. *Pax6*-dependence of *Six3*, *Eya1* and *Dach1* expression during lens and nasal placode induction. *Gene Expr. Patterns* 6, 110–118.
- Quiring, R., Walldorf, U., Kloter, U., Gehring, W.J., 1994. Homology of the eyeless gene of *Drosophila* to the small eye gene in mice and Aniridia in humans. *Science* 265, 785–789.
- Rayapureddi, J.P., Kattamuri, C., Steinmetz, B.D., Frankfort, B.J., Ostrin, E.J., Mardon, G., Hegde, R.S., 2003. Eyes absent represents a class of protein tyrosine phosphatases. *Nature* 426, 295–298.
- Sahly, I., Andermann, P., Petit, C., 1999. The zebrafish *eya1* gene and its expression pattern during embryogenesis. *Dev. Genes Evol.* 209, 399–410.
- Salzer, C.L., Kumar, J.P., 2009. Position dependent responses to discontinuities in the retinal determination network. *Dev. Biol.* 326, 121–130.
- Schlosser, G., 2006. Induction and specification of cranial placodes. *Dev. Biol.* 294, 303–351.
- Schlosser, G., Ahrens, K., 2004. Molecular anatomy of placode development in *Xenopus laevis*. *Dev. Biol.* 271, 439–466.

- Schlosser, G., Awtry, T., Brugmann, S.A., Jensen, E.D., Neilson, K., Ruan, G., Stammier, A., Voelker, D., Yan, B., Zhang, C., Klymkowsky, M.W., Moody, S.A., 2008. Eya1 and Six1 promote neurogenesis in the cranial placodes in a SoxB1-dependent fashion. *Dev. Biol.* 320, 199–214.
- Seimiya, M., Gehring, W.J., 2000. The *Drosophila* homeobox gene optix is capable of inducing ectopic eyes by an eyeless-independent mechanism. *Development* 127, 1879–1886.
- Serikaku, M.A., O'Tousa, J.E., 1994. sine oculis is a homeobox gene required for *Drosophila* visual system development. *Genetics* 138, 1137–1150.
- Sheng, G., Stern, C.D., 1999. Gata2 and Gata3: novel markers for early embryonic polarity and for non-neural ectoderm in the chick embryo. *Mech. Dev.* 87, 213–216.
- Shen, W., Mardon, G., 1997. Ectopic eye development in *Drosophila* induced by directed dachshund expression. *Development* 124, 45–52.
- Silver, S.J., Rebay, I., 2005. Signaling circuitries in development: insights from the retinal determination gene network. *Development* 132, 3–13.
- Solomon, K.S., Fritz, A., 2002. Concerted action of two dlx paralogs in sensory placode formation. *Development* 129, 3127–3136.
- Spitz, F., Demignon, J., Porteu, A., Kahn, A., Concordet, J.P., Daegelen, D., Maire, P., 1998. Expression of myogenin during embryogenesis is controlled by six/sine oculis homeoproteins through a conserved mef3 binding site. *Proc. Natl. Acad. Sci. U. S. A.* 95, 14220–14225.
- Stark, M.R., Sechrist, J., Bronner-Fraser, M., Marcelle, C., 1997. Neural tube-ectoderm interactions are required for trigeminal placode formation. *Development* 124, 4287–4295.
- Stern, C.D., Ireland, G.W., 1981. An integrated experimental study of endoderm formation in avian embryos. *Anat. Embryol. (Berl.)* 163, 245–263.
- Streit, A., 2002. Extensive cell movements accompany formation of the otic placode. *Dev. Biol.* 249, 237–254.
- Streit, A., 2004. Early development of the cranial sensory nervous system: from a common field to individual placodes. *Dev. Biol.* 276, 1–15.
- Streit, A., 2007. The preplacodal region: an ectodermal domain with multipotential progenitors that contribute to sense organs and cranial sensory ganglia. *Int. J. Dev. Biol.* 51, 447–461.
- Streit, A., Lee, K.J., Woo, I., Roberts, C., Jessell, T.M., Stern, C.D., 1998. Chordin regulates primitive streak development and the stability of induced neural cells, but is not sufficient for neural induction in the chick embryo. *Development* 125, 507–519.
- Suzuki-Yagawa, Y., Kawakami, K., Nagano, K., 1992. Housekeeping Na, K-ATPase alpha 1 subunit gene promoter is composed of multiple cis elements to which common and cell type-specific factors bind. *Mol. Cell. Biol.* 12, 4046–4055.
- Tootle, T.L., Silver, S.J., Davies, E.L., Newman, V., Latek, R.R., Mills, I.A., Selengut, J.D., Parlikar, B.E., Rebay, I., 2003. The transcription factor Eyes absent is a protein tyrosine phosphatase. *Nature* 426, 299–302.
- Torres, M., Gomez-Pardo, E., Gruss, P., 1996. Pax2 contributes to inner ear patterning and optic nerve trajectory. *Development* 122, 3381–3391.
- Treisman, J.E., 1999. A conserved blueprint for the eye? *Bioessays* 21, 843–850.
- Uwanogho, D., Rex, M., Cartwright, E.J., Pearl, G., Healy, C., Scotting, P.J., Sharpe, P.T., 1995. Embryonic expression of the chicken Sox2, Sox3 and Sox11 genes suggests an interactive role in neuronal development. *Mech. Dev.* 49, 23–36.
- Wawersik, S., Maas, R.L., 2000. Vertebrate eye development as modeled in *Drosophila*. *Hum. Mol. Genet.* 9, 917–925.
- Weasner, B., Salzer, C., Kumar, J.P., 2007. Sine oculis, a member of the SIX family of transcription factors, directs eye formation. *Dev. Biol.* 303, 756–771.
- Woda, J.M., Pastagia, J., Mercola, M., Artinger, K.B., 2003. Dlx proteins position the neural plate border and determine adjacent cell fates. *Development* 130, 331–342.
- Xu, H., Dude, C.M., Baker, C.V.H., 2008. Fine-grained fate maps for the ophthalmic and maxillomandibular trigeminal placodes in the chick embryo. *Dev. Biol.* 317, 174–186.
- Xu, P.X., Adams, J., Peters, H., Brown, M.C., Heaney, S., Maas, R., 1999. Eya1-deficient mice lack ears and kidneys and show abnormal apoptosis of organ primordia. *Nat. Genet.* 23, 113–117.
- Zheng, W., Huang, L., Wei, Z.B., Silvius, D., Tang, B., Xu, P.X., 2003. The role of Six1 in mammalian auditory system development. *Development* 130, 3989–4000.
- Zou, D., Silvius, D., Fritzsche, B., Xu, P.X., 2004. Eya1 and Six1 are essential for early steps of sensory neurogenesis in mammalian cranial placodes. *Development* 131, 5561–5572.
- Zou, D., Silvius, D., Rodrigo-Blomqvist, S., Enerback, S., Xu, P.-X., 2006. Eya1 regulates the growth of otic epithelium and interacts with Pax2 during the development of all sensory areas in the inner ear. *Dev. Biol.* 298, 430–441.